



## General production advice ware potatoes

- \* **High Yields**
- \* **Bold Tubers**
- \* **Excellent Quality**
- \* **Multiuse**
- \* **Suitable For A Range Of Soil Types**
- \* **Good Storage Characteristics**



<b>Maturity</b>	<b>Second Early</b>
<b>Tuber Size</b>	<b>Large</b>
<b>Tuber Shape</b>	<b>Oval</b>
<b>Numbers of Tubers</b>	<b>8-10</b>
<b>Flesh Colour</b>	<b>Yellow</b>
<b>Skin Colour</b>	<b>Yellow</b>
<b>Berries</b>	<b>No Berries</b>
<b>Dormancy</b>	<b>Medium</b>
<b>Emergence</b>	<b>Normal</b>
<b>Sencorex Sensitivity</b>	<b>Pre Emergence only</b>
<b>Foliage Development</b>	<b>Vigorous</b>
<b>Dry Matter</b>	<b>21.7%</b>
<b>Foliage Blight</b>	<b>6</b>
<b>Tuber Blight</b>	<b>7</b>
<b>Blackleg</b>	<b>6.5</b>
<b>Common Scab</b>	<b>5.5</b>
<b>Powdery Scab</b>	<b>6.5</b>
<b>Gangrene</b>	<b>-</b>
<b>Potato Leaf Roll Virus</b>	<b>-</b>
<b>Potato Virus Y</b>	<b>-</b>
<b>PVYn</b>	<b>-</b>
<b>Yntn Tuber Tolerance</b>	<b>Resistant</b>
<b>Spraing</b>	<b>8</b>
<b>Black Scurf</b>	<b>5</b>
<b>Dry Rot</b>	<b>7</b>
<b>Slugs</b>	<b>5</b>
<b>Internal Bruising</b>	<b>3 (1=Low, 20=Sensitive)</b>
<b>Little Potato Disorder</b>	<b>4</b>
<b>PCN R01,4</b>	<b>Resistant</b>
<b>PCN Pa2 Pa3</b>	<b>Susceptible</b>

**(1=Susceptible, 9=Resistant)**




### Fertiliser

**Sagitta has a high to moderate Nitrogen requirement, Advice 200-250 kg/ha inclusive of soil supply. 2/3 before planting 1/3 directly after tuberisation.**

**Phosphate and Potassium according to RB209**

## General production advice ware potatoes

### Pre-treatment and planting




Sagitta is susceptible to Little Potato Disorder, manage the seed accordingly to minimise excessive temperature fluctuation and uncontrolled sprouting. Treatment against Rhizoctonia is recommended. A good formation of the ridge is recommended to help prevent greening.

### Growing attention points




Sagitta is susceptible to Alternaria, preventative treatment is recommended. Choose light to medium soil types to obtain the best tuber numbers and shape, the variety has a slight susceptibility to second growth.

### Haulm killing and harvest



Sagitta can bulk quickly, therefore care is required with the timing of desiccation.

### Storage



Sagitta does not have a long dormancy therefore for longer term storage the first application of sprout inhibitors should be made in good time.